

Raman Scattering from Frustrated Quantum Spin Chains

Wolfram Brenig

Institut für Theoretische Physik, Universität zu Köln Zùlpicher Str. 77, 50937 Köln, Germany

(February 1, 2008)

Abstract

Magnetic Raman scattering from a frustrated spin-1/2 Heisenberg-chain is considered with a focus on the uniform phase of the spin-Peierls compound CuGeO_3 . The Raman intensity is analyzed in terms of a Loudon-Fleury scattering process using a spinless-fermion mean-field theory developed for the frustrated spin-chain. A comparison to experimental data is presented and the frustration and temperature dependence is studied. In good agreement with observed spectra a broad inelastic four-spinon continuum is found at low temperatures above the spin-Peierls transition. At high temperatures the intensity develops a quasi-elastic line analogous to experiment.

PACS: 75.10.Jm, 75.40.Gb, 78.30.-j

I. INTRODUCTION

Elementary excitations of one-dimensional (1D) quantum spin-chains exhibit a number of remarkable subtleties, such as the magnon continuum of the isotropic 1D spin-1/2 Heisenberg model or the excitation-gap for integer-spin systems as conjectured by Haldane. Recent and extensive studies of 'spin-ladder' compounds as well as of the germanate CuGeO_3 [1] have lead to renewed interest in quantum spin-chains. In this context, magnetic Raman scattering, if symmetry-allowed, has developed into an important tool to investigate the local spin dynamics [2–4]. Here I will establish a simple framework to interpret the magnetic Raman scattering from a frustrated quantum spin-chain with a particular focus on the uniform phase of CuGeO_3 .

CuGeO_3 is a quasi-1D anorganic spin-Peierls compound with a dimerization transition at a temperature $T_{SP} \simeq 14K$ [1,5,6]. Its structure comprises of weakly coupled CuO_2 chains, with copper in a spin-1/2 state [7]. Since the nearest-neighbor ($n.n.$) exchange-coupling between copper spins along the CuO_2 chains is strongly reduced by almost orthogonal intermediate oxygen states [8] the next-nearest-neighbor ($n.n.n.$) exchange is relevant. Both, $n.n.$ and $n.n.n.$ exchange, are antiferromagnetic [9] implying intra-chain frustration. A minimal model of CuGeO_3 is the J_1 - J_2 - δ model

$$H = J_1 \sum_l [(1 + (-)^l \delta) \mathbf{S}_l \cdot \mathbf{S}_{l+1} + \alpha \mathbf{S}_l \cdot \mathbf{S}_{l+2}] \quad . \quad (1)$$

Here, $\mathbf{S}_l$ is a spin-1/2 operator, $J_1 \approx 160K$ [10–12] is the $n.n.$ exchange-coupling constant and δ resembles the lattice dimerization which is finite for $T < T_{SP}$ only. α is the intra-chain frustration-ratio $\alpha = J_2/J_1$ where J_2 is the $n.n.n.$ exchange-coupling constant. A final consensus on the precise magnitude of α is still lacking. Studies of the magnetic susceptibility have resulted in $\alpha \approx 0.24$ [10] as well as in $\alpha \approx 0.35$ [11]. The latter is consistent with a comparative investigation of magnetic susceptibility *and* thermal expansion [12]. Therefore, very likely, CuGeO_3 displays a frustration induced contribution to the spin-gap at zero temperature irrespective of the actual lattice dimerization [13]. Depending on

the magnitude of the frustration the zero-temperature dimerization $\delta(T = 0)$ has to be on the order of $\delta(0) = 0.01...0.03$ such as to enforce the size of the spin-gap observed in inelastic neutron scattering (INS) [5].

Magnetic excitations in CuGeO_3 are clearly distinct among the uniform, i.e. $T > T_{SP}$, and the dimerized, i.e. $T < T_{SP}$, phase. While the dynamic structure factor exhibits a gapless two-spinon continuum similar to that of the 1D Heisenberg chain above T_{SP} [14], well defined magnon-like excitations have been observed below T_{SP} [15]. These magnons reside within the spin-gap and are split off from the two-spinon continuum. They have been interpreted as two-spinon bound states [16,17].

Magnetic Raman scattering from CuGeO_3 has been observed both, in the dimerized as well as in the uniform phase [2–4]. In the low temperature uniform phase, for $T_{SP} < T \ll J_1$, the Raman spectrum displays a broad continuum centered at $\hbar\omega \approx 2.4J_1$. Early on this continuum has been related to four-spinon excitations [2–4,18]. In the high temperature uniform phase, and in addition to the four-spinon continuum, the spectra show a pronounced quasi-elastic line. In the dimerized phase the Raman intensity develops a gap at approximately $1.5\text{--}1.8\Delta_{ST}$ [20] where Δ_{ST} is the singlet-triplet gap observed in INS [5]. Moreover, four characteristic peaks appear in the spectrum the lowest one of which at 30cm^{-1} ‘coincides’ with the Raman gap. At present the interpretations of these peaks are controversial. Tentatively the 30cm^{-1} line has been attributed to a *continuum* of two-magnon bound states [19,20]. A possible dimensional crossover effect has been invoked for the 225cm^{-1} line [3,20]. In this respect the influence of inter-chain coupling in CuGeO_3 is an open problem [21].

Here I will focus on Raman scattering from the uniform phase of the $J_1\text{--}J_2\text{--}\delta$ model, i.e. at $\delta = 0$. First I will describe a spinless-fermion mean-field theory to treat the $J_1\text{--}J_2$ Hamiltonian. Next the Raman intensity is expressed in terms of a four-fermion correlation function. Finally results for the Raman spectra are compared with experimental findings and are contrasted against other theoretical approaches.

II. MEAN-FIELD THEORY OF THE J_1 - J_2 MODEL

The spinless-fermion mean-field (MF) theory for the J_1 - J_2 model is based on the Jordan-Wigner (JW) representation [22] of the spin algebra, i.e. $S_l^z = (c_l^\dagger c_l - \frac{1}{2})$ and $S_l^+ = (-)^l \prod_{j < l} (1 - c_j^\dagger c_j) c_l^\dagger$, where $c_l^{(\dagger)}$ are fermion operators. Inserting this into (1) at $\delta = 0$ one obtains

$$H = \sum_l \left[-\frac{1}{2} c_l^\dagger c_{l+1} - \frac{1}{2} c_{l+1}^\dagger c_l - c_l^\dagger c_l + c_l^\dagger c_l c_{l+1}^\dagger c_{l+1} + \frac{1}{4} + \frac{\alpha}{2} c_l^\dagger c_{l+2} + \frac{\alpha}{2} c_{l+2}^\dagger c_l \right. \\ \left. + \alpha (-c_l^\dagger c_{l+2} c_{l+1}^\dagger c_{l+1} - c_{l+1}^\dagger c_{l+1} c_{l+2}^\dagger c_l - c_l^\dagger c_l + c_l^\dagger c_l c_{l+2}^\dagger c_{l+2} + \frac{1}{4}) \right] , \quad (2)$$

where J_1 is set to unity in section II and III. l runs over the lattice sites. The terms proportional to α are absent in the JW representation of the isotropic $n.n.$ Heisenberg model. In contrast to the $n.n.$ model, both, the transverse as well as the longitudinal $n.n.n.$ exchange-interaction lead to four-fermion vertices. Longer range spin-exchange leads to even higher order couplings, as is obvious from the JW representation which implies $2l$ -fermion vertices for l -th.-nearest-neighbor spin-exchange. To proceed I treat Hamiltonian (2) in MF-approximation. Allowing for all contractions of type $\langle c_l^\dagger c_m \rangle$ one gets

$$H_{MF} = \sum_k \left\{ -[1 + (A + B)(1 - 2\alpha)] \cos(k) c_k^\dagger c_k \right. \\ \left. + \frac{i}{2} (A - B)(1 + 2\alpha) \sin(k) (c_k^\dagger c_{k+\pi} - c_{k+\pi}^\dagger c_k) \right\} + const. , \quad (3)$$

where k is the momentum, $A = A^* = \langle c_{2l}^\dagger c_{2l+1} \rangle$, and $B = B^* = \langle c_{2l+1}^\dagger c_{2l} \rangle$. In principle contractions of type $D = D^* = \langle c_{2l+2}^\dagger c_{2l} \rangle$ do occur, however, their selfconsistent value can be shown to vanish identically. Note that (3) allows for both, a uniform, if $A - B = 0$, and a gaped phase, if $A - B \neq 0$. Here no attempt will be made to describe the spin-dimer state using $A \neq B$ and the relevance of the gaped solution of the MF-theory will be discussed elsewhere. In the uniform case (3) resembles a single-band spinon gas with a hopping amplitude $t(T, \alpha)$ to be determined selfconsistently

$$H_{MF} = \sum_k \epsilon_k c_k^\dagger c_k$$

$$\epsilon_k = -t(T, \alpha) \cos(k) = -[1 + 2A(1 - 2\alpha)] \cos(k) \quad (4)$$

$$A = \frac{2}{N} \sum_{0 \leq k \leq \pi} \cos(k) f(\epsilon_k) \quad ,$$

where $f(\epsilon) = [\exp(\epsilon/T) + 1]^{-1}$ is the Fermi function. At zero temperature the spinon dispersion simplifies to

$$\epsilon_k(T = 0) = -[1 + 2(1 - 2\alpha)/\pi] \cos(k) \quad . \quad (5)$$

For vanishing frustration this is identical to Bulaevskii's result $-(1 + 2/\pi) \cos(k)$ [23]. The latter is known to compare reasonably well with the exact spinon dispersion $\epsilon_k = -\pi/2 \cos(k)$ by des Cloizeaux and Pearson [24]. For finite α the MF-theory results in a frustration induced *softening* of the spinon stiffness which, for $T = 0$, can be expressed as

$$\frac{v_s(\alpha)}{v_s(0)} = 1 - \frac{4}{2 + \pi} \alpha \approx 1 - 0.778\alpha \quad , \quad (6)$$

where $v_s(\alpha) = \partial \epsilon_k / \partial k|_{k=\pi/2}$ is the spinon velocity. Eqn. (6) is qualitatively consistent with a numerical study where $v_s(\alpha)/v_s(0) \approx 1 - 1.12\alpha$ has been found [17]. In fig. 1 $t(T, \alpha)$ is depicted as a function of temperature for various values of frustration. As is obvious the MF spinon-stiffness is decreased, both, as a function of increasing frustration and temperature. At zero temperature the MF ground state energy is given by $E_0(\alpha) = N(-1/\pi - 1/\pi^2 + 2\alpha/\pi^2)$ which, for $\alpha = 0$, leads to $E_0(0) \approx -0.420$ [23]. This agrees reasonably well with the Bethe-Ansatz result $E_0 = 1/4 - \ln(2) \approx -0.443$. Moreover, the linear frustration dependence of the MF ground state energy, $(E_0(\alpha) - E_0(0))/N \approx 0.203\alpha$ is close to that found in finite-chain diagonalization [25], where $(E_0(\alpha) - E_0(0))/N \approx 0.177\alpha$ for $\alpha \lesssim 0.3$.

III. RAMAN SCATTERING

The MF-theory is a convenient tool to study Raman scattering from the J_1 - J_2 model. The Raman vertex is given by Loudon-Fleury's photon-induced super-exchange operator [26]

$$R = \sum_{lm} T_{lm} (\mathbf{E}_{in} \cdot \mathbf{n}_{lm}) (\mathbf{E}_{out} \cdot \mathbf{n}_{lm}) \mathbf{S}_l \cdot \mathbf{S}_m \quad . \quad (7)$$

Here T_{lm} sets the coupling strength, $\mathbf{E}_{in}$ ($\mathbf{E}_{out}$) refers to the field of the in(out)going light, and $\mathbf{n}_{lm}$ labels a unit vector connecting the sites l and m . By symmetry, in a strictly 1D situation, (7) leads to scattering only for parallel polarization along the c -directed chains. Since the real-space decay of T_{lm} is comparable to that of the exchange integrals in (1) it is sufficient to consider a Raman operator with at most *n.n.n.* spin-exchange

$$R = C \sum_l (\mathbf{S}_l \cdot \mathbf{S}_{l+1} + \beta \mathbf{S}_l \cdot \mathbf{S}_{l+2}) \quad . \quad (8)$$

The scattering intensity $I(\omega)$ is obtained via the fluctuation dissipation theorem $I(\omega) = \chi''(\omega)/(1 - e^{-\omega/T})$ from the dynamical susceptibility of the Raman operator

$$\chi''(\omega) = \text{Im}[\chi(\omega + i\eta)] = \text{Re} \int_0^{+\infty} dt e^{i(\omega + i\eta)t} \langle [R(t), R] \rangle \quad , \quad (9)$$

where $\langle \dots \rangle$ denotes the thermal average.

Frustration of *unequal* magnitude regarding the J_1 - J_2 model and the Raman operator is *mandatory* for non-vanishing inelastic scattering. If $\alpha = \beta$ the Hamiltonian for $\delta = 0$ and the Raman operator commute which leads to elastic scattering only. The inelastic intensity which results from (8) is identical to that of a Raman operator $R' = R - \gamma H$. Setting $\gamma = C\beta/\alpha$ or $\gamma = C$ leads to scattering by a renormalized *n.n.* or *n.n.n.* Raman operator only

$$R_1 = C(1 - \frac{\beta}{\alpha}) \sum_l \mathbf{S}_l \cdot \mathbf{S}_{l+1} \quad \text{or} \quad R_2 = C(\beta - \alpha) \sum_l \mathbf{S}_l \cdot \mathbf{S}_{l+2} \quad . \quad (10)$$

At present, an *exact* treatment of (9) is not feasible and the Raman intensity of any *approximate* evaluation will depend on γ . In this respect R_2 is the proper choice for $\alpha, \beta \ll 1$ since it guarantees that $I(\omega) \propto (\beta - \alpha)^2$. This is not obvious for R_1 or other values of γ . In those cases, even for $\alpha, \beta \rightarrow 0$, R' may contain a component of order unity, proportional to the Hamiltonian, which has to be projected out. To avoid this complication R_2 will be considered hereafter.

Using the JW-fermions the Raman operator R_2 can be expressed in terms of a four-fermion operator

$$R_2 = C(\beta - \alpha) \sum_{k,k',q} h(k, k', q) c_k^\dagger c_{k+q} c_{k'}^\dagger c_{k'-q} + \Lambda_{1ph} \quad (11)$$

$$h(k, k', q) = \cos(2q) - \cos(2k + q) - \cos(2k' - q) \quad ,$$

where Λ_{1ph} labels all one-particle-hole excitations of the JW-transform of R_2 . Since these occur at zero total momentum they do not contribute to inelastic Raman scattering within MF-theory [27]. Using (11), the Raman intensity (9) is written in terms of the four-fermion propagator

$$\chi(\tau) = C^2(\alpha - \beta)^2 \sum_{k,k',q,p,p',r} [h(k, k', q)h(p, p', r) \langle T_\tau(c_k^\dagger(\tau)c_{k+q}(\tau)c_{k'}^\dagger(\tau)c_{k'-q}(\tau)c_{p'-r}^\dagger c_{p'+r}^\dagger c_p) \rangle] \quad . \quad (12)$$

Here τ is the imaginary time and $\chi''(\omega)$ results from the usual analytic continuation $\chi''(\omega) = -Im[\chi(i\omega_n \rightarrow \omega + i0^+)]$ where $\omega_n = 2n\pi T$ is a Bose Matsubara-frequency. On the level of MF-theory the four-fermion propagator (12) is evaluated neglecting all vertex corrections and using the MF one-particle Green's functions corresponding to (4). After a number of standard manipulations I obtain

$$\chi''_{MF}(\omega) = -\frac{1}{2\pi^2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dk \sum_{k'} \left\{ \frac{g^2(k, k' + q, q)}{\sqrt{[2t \sin(q/2)]^2 - (\epsilon_{k+q} - \epsilon_k - \omega)^2}} [f(\epsilon_k) - f(\epsilon_{k+q})][f(\epsilon_{k'}) - f(\epsilon_{k'+q})][n(\epsilon_{k'+q} - \epsilon_{k'} + \omega) - n(\epsilon_{k'+q} - \epsilon_{k'})] \right\} \quad , \quad (13)$$

where $g(k, k', q) = \frac{1}{2}[h(k, k', q) + h(k, k', k' - k - q)]$ and the discrete sum on k' runs over the set of solutions of

$$\sin(k' + q/2) = \frac{\epsilon_{k+q} - \epsilon_k - \omega}{2t \sin(q/2)} \quad ; \quad k' \in] - \pi, \pi] \quad . \quad (14)$$

This concludes the MF-theory of Raman scattering.

IV. DISCUSSION

In fig. 2 I compare the MF-intensity obtained from a numerical integration of (13) (solid and dashed lines) at $T = 20K$ and $12.8K$, for $J_1 = 160K$ and two values of frustration $\alpha = 0.24$ and $\alpha = 0.35$ with an experimental Raman spectrum observed at $T = 20K$ in the uniform phase of CuGeO_3 [4,18] (dashed-dotted line with markers). Phonon lines at 184cm^{-1} and 330cm^{-1} have been removed. This Raman data is consistent with other published results [2,3]. The absolute magnitude of the observed Raman intensity as well as the coupling strength C of the scattering operator (8) are unknown quantities. Therefore the two sets of MF-intensities for distinct α and the experimental spectrum have been normalized on a scale of arbitrary units. The theoretical parameters in fig. 2 represent no attempt at a 'best fit' to the data, rather they have been chosen among those obtained independently from studies of the magnetic susceptibility of CuGeO_3 [10–12]. Figure 2 displays reasonable agreement between theory and experiment, although details of the experimental spectrum, i.e. the structure at 375 wave numbers and the initial curvature are not reproduced by the MF-theory. Evidently a value of $\alpha = 0.24$ [10] is slightly more favored by this comparison at $J_1 = 160K$ than $\alpha = 0.35$ [11,12]. Similar findings have been made in numerical studies [20]. However, since the maximum of the MF-intensity roughly scales with the maximum of the spinon-dispersion $\propto J_1[1 + 2(1 - 2\alpha)/\pi]$, a better agreement with $\alpha = 0.35$ instead of $\alpha = 0.24$ can also be reached simply by increasing J_1 by $\sim 10\%$.

The comparison in fig. 2 strongly corroborates a study [18] of Raman scattering from the uniform phase of CuGeO_3 which is based the 'solitonic' mean-field description of the XXZ Heisenberg chain by Gómez-Santos [28]. This approach employs a domain-wall representation of the spin chain which leads to formal developments very different from the straightforward application of the Jordan-Wigner type of MF-theory presented here.

In fig. 3 the temperature dependence of the MF-spectrum is shown, both, in terms of the intensity $I(\omega)$ – which is observed in experiment – and the Raman-operator susceptibility $\chi''(\omega)$. The latter exhibits a left-shift of its maximum upon increase of the temperature.

This leads to enhanced low-frequency spectral weight which, by virtue of the Bose-prefactor, turns into a quasi-elastic line in $I(\omega)$ for $T \gtrsim J_1$. A corresponding tendency has been detected in finite-temperature Lanczos studies [32]. In MF-theory the high-temperature enhancement of the low-frequency spectral weight in $\chi''(\omega)$ is due to the reduction of the spinon-stiffness as a function of increasing temperature, see fig. 1, and due to the 'smearing' induced by the Fermi- and Bose-functions contained in (13). It is tempting to relate the behavior depicted in fig. 3 to the high-temperature quasi-elastic line observed in the uniform phase of CuGeO_3 [2–4]. However, in CuGeO_3 the four-spinon continuum is only weakly shifted by temperature and remains more clearly separated from the quasi-elastic line for $T_{SP} \lesssim T \lesssim J_1$. This discrepancy may be due to an overestimation of the temperature dependence of the spinon-stiffness in MF-theory [29,30].

Finally, I emphasize that the MF-theory gives only a limited description of spinon interaction effects. In the case of the *two*-spinon propagator this is known to result in an incorrect description of the spectral weight distribution [31]. This caveat of MF-theory has stimulated a study of approximate vertex corrections to the *four*-spinon-propagator [32]. However, the resulting Raman spectra show no agreement with experiment. At present the impact of spinon interaction effects beyond MF-theory on the Raman spectra remain unclear.

In conclusion I have described a finite temperature MF-theory for frustrated spin-chains and consequently detailed its application to magnetic Raman scattering. At low-temperatures, in the gapless phase, I find a scattering continuum which is due to frustration induced four-spinon excitations and is compatible with observed Raman spectra of CuGeO_3 . In the high-temperature regime the MF Raman-intensity is dominated by a quasi-elastic line which results from the temperature dependence of the spinon spectrum.

V. ACKNOWLEDGMENTS

I am indebted to B. Büchner, P. Knoll, and E. Müller–Hartmann for valuable discussions. Part of this work has been supported by the Deutsche Forschungsgemeinschaft through the SFB 341.

REFERENCES

- [1] M. Hase *et al.*, Phys. Rev. Lett. **70**, 3651 (1993).
- [2] H. Kuroe *et al.*, Phys. Rev. B **50**, 16468 (1994).
- [3] P.H.M. van Loosdrecht *et al.*, Phys. Rev. Lett. **76**, 311 (1996).
- [4] P. Lemmens *et al.*, Physica B **223&224**, 535 (1996).
- [5] M. Nishi *et al.*, Phys. Rev. B **50**, 6508 (1994); M.C. Martin *et al.*, Phys. Rev. B **53**, R14713 (1996). L.P. Regnault *et al.*, Physica B **213 & 214**, 278 (1995);
- [6] J.P. Pouget *et al.*, Phys. Rev. Lett. **72**, 4037 (1994); O. Kamimura *et al.*, J. Phys. Soc. Jpn. **63**, 2467 (1994); S.B. Oseroff *et al.*, Phys. Rev. Lett. **74**, 1450 (1995).
- [7] H. Völlenkle *et al.*, Monatsh. Chem. **98**, 1352 (1967); G.A. Petrakovskii *et al.*, Zh. Eksp. Teor. Fiz. **98**, 1382 (1990) [Sov. Phys. JETP **71**, 772 (1990)]; K. Hirota *et al.*, Phys. Rev. Lett. **73**, 736 (1994); J.E. Lorenzo *et al.*, Phys. Rev. B **50**, 1278 (1994); Q.J. Harris *et al.*, Phys. Rev. B **50**, 12606 (1994); M. Arai *et al.*, J. Phys. Soc. Jpn. **63**, 1661 (1994).
- [8] M. Braden *et al.*, Phys. Rev. B **54**, 1105 (1996).
- [9] W. Geertsma and D. Khomskii, Phys. Rev. B **54**, 3011 (1996).
- [10] G. Castillia *et al.*, Phys. Rev. Lett. **75**, 1823 (1995).
- [11] J. Riera *et al.*, Phys. Rev. B **51**, 16098 (1995).
- [12] B. Büchner, unpublished.
- [13] R. Chitra *et al.*, Phys. Rev. B **52**, 6581 (1995).
- [14] M. Arai *et al.*, Phys. Rev. Lett. **77**, 3649 (1996).
- [15] M. Aín *et al.*, preprint.
- [16] G.S. Uhrig and H.J. Schulz, Phys. Rev. B **54**, R9624 (1996).

- [17] A. Fledderjohann and C. Gros, preprint cond-mat/9612013
- [18] V.N. Muthukumar *et al.*, Phys. Rev. B **54**, R9635 (1996).
- [19] P.H.M. van Loosdrecht *et al.*, preprint cond-mat/9612167.
- [20] C. Gros *et al.*, preprint cond-mat/9612101.
- [21] G.S. Uhrig, preprint, (1996).
- [22] P. Jordan and E. Wigner, Z. Phys. **47**, 631 (1928).
- [23] L.N. Bulaevskii *et al.*, Zh. Eksp. Teor. Fiz. **43**, 968 (1962), [Sov. Phys. JETP **16**, 685 (1963)]
- [24] J. des Cloizeaux and J.J. Pearson, Phys. Rev. **128**, 2131 (1962).
- [25] T. Tonegawa and I. Harada, J. Phys. Soc. Jpn. **56**, 2153 (1987).
- [26] P.A. Fleury and R. Loudon, Phys. Rev. **166**, 514 (1968).
- [27] In Raman scattering photon-momentum transfer to the electronic system can be neglected which is implicit in the definition of R . Therefore spinon one-particle-hole excitations will contribute to the inelastic intensity only in case of a finite quasi-particle relaxation rate, which is absent in MF-theory.
- [28] G. Gómez-Santos, Phys. Rev. B **41**, 6788 (1990).
- [29] K. Frabricsius *et al.*, preprint cond-mat/9611077
- [30] O. A. Starykh *et al.*, preprint cond-mat/9701052.
- [31] G. Müller, H. Beck, and J.C. Bonner, Phys. Rev. Lett. **43**, 75 (1979).
- [32] R.P.R. Singh *et al.*, preprint (1996).

FIGURES

FIG. 1. Temperature dependence of the mean-field hopping amplitude for various values of frustration.

FIG. 2. Mean-field Raman intensity (solid and dashed lines) at two temperatures and for two values of frustration as compared to the experimental Raman spectrum of CuGeO_3 [4,18] (dashed-dotted line & markers).

FIG. 3. Mean-field Raman intensity and susceptibility for various temperatures.




Fig. 1, Brenig




Fig. 2, Brenig




Fig. 3, Brenig